



Instructions for Use
Medication: Tri-Caine Blue Anesthetic Gel
Lidocaine 10%, Prilocaine 10%, Tetracaine 4%,

Please Note:

Maximum recommended adult dose is 0.5mls and 0.2mls in children weighing 40 pounds or more.

Patients should not eat or drink anything until full sensation returns to lips, mouth, and throat. Use caution with heat or cold; Patient will not have accurate hot or cold sensation until full effects of anesthesia have worn off.

Directions

1. Dry mucosal area.
2. Apply a regular sized micro-brush amount to dry, oral mucosa and leave in, with applicator in place, for 1-3 minutes. This will allow for the medication to be absorbed and have a lasting anesthetic effect.
3. Rinse area to remove left over gel prior to starting procedure. (Peak anesthetic effect at approximately 6 minutes; anesthetic effect lasts approximately 20-30 minutes.)

If using medication over a large area or quadrant:

1. Dry mucosal area.
2. Apply two to three times the amount of medication (not to exceed the maximum dose) with a micro-brush and smear it thoroughly over quadrant/area for a period of 1 minute. Keep area as dry as possible for best mucosal absorption and for longer duration of anesthetic effect.
3. Wait until area is numb prior to beginning procedure (3-6 minutes).
4. Rinse area to remove left over gel prior to starting procedure. (Peak anesthetic effect at approximately 6 minutes; anesthetic effect lasts approximately 20-30 minutes.)